

macular pucker


a closer look

HOW THE EYE SEES

The **macula** is the small area at the center of the eye's **retina** that allows you to see fine details clearly. The retina is a layer of light-sensing cells lining the back of your eye.

As light rays enter your eye, the retina converts the rays into signals, which are sent through the optic nerve to your brain where they are recognized as images.

Damage to your macula causes blurred central vision, making it difficult to perform tasks such as reading small print or threading a needle.



WHAT IS A MACULAR PUCKER?

The macula normally lies flat against the back of the eye, like film lining the back of a camera. When wrinkles, creases or bulges form on the macula, this is known as **macular pucker**.

Your ophthalmologist uses tiny instruments to remove the wrinkled tissue on the macula.

WHAT ARE THE SYMPTOMS OF MACULAR PUCKER?

Symptoms of macular pucker range from mild to severe and may involve one or both eyes.

Symptoms may include:

- Blurred central (detail) vision;
- Distorted, or "wavy," vision;
- Difficulty reading or performing tasks that require detail vision;
- Gray and/or cloudy area in central vision;
- Central blind spot.

Peripheral (side) vision is not affected.



WHAT CAUSES MACULAR PUCKER?

As you age, the vitreous — the clear, gel-like substance that fills the middle of your eye — begins to shrink and pull away from the retina. As the vitreous pulls away, scar tissue may develop on the macula. Sometimes the scar tissue can warp and contract, causing the retina to wrinkle or bulge.

Eye conditions associated with macular pucker include:

- Vitreous detachment;
- Torn or detached retina;
- Inflammation inside the eye;
- Severe trauma to the eye (from surgery or injury);
- Disorders of the blood vessels in the retina.

HOW IS MACULAR PUCKER DETECTED?

Your ophthalmologist (Eye M.D.) detects macular pucker by examining your retina. Your doctor may perform fluorescein angiography or optical coherence tomography (OCT) — procedures that take special photographs of the eye. These photographs show if an abnormality exists in your retina.

HOW IS MACULAR PUCKER TREATED?

For mild symptoms, no treatment may be necessary. Updating your eyeglass prescription or wearing bifocals may improve vision. Eyedrops, medicines or laser surgery do not improve vision.

For more severe symptoms, a surgery called **vitrectomy** is recommended. The surgery is usually performed as an outpatient procedure in an operating room. During surgery, your ophthalmologist uses tiny instruments to remove the wrinkled tissue on your macula. After the tissue is gone, the macula flattens and vision slowly improves, though it usually does not return all the way to normal.

You should consider surgery if your blurred vision is interfering with your daily activities.

ARE THERE ANY RISKS INVOLVED WITH VITRECTOMY SURGERY?

As with any surgical procedure, complications can occur with vitrectomy surgery. Complications may include:

- Infection;
- Bleeding;
- Retinal detachment;
- Recurrence of macular pucker.

After vitrectomy surgery, cataracts (clouding of the eye's lens) may also develop. Be sure to discuss potential complications with your ophthalmologist before surgery.

macular pucker



NOTES

COMPLIMENTS OF YOUR OPHTHALMOLOGIST:

San Antonio Eye Center, P.A.
(210) 226-6169
www.saeye.com

Academy reviewed 03/11

© 2011 American Academy of Ophthalmology. The American Academy of Ophthalmology, The Eye M.D. Association and the Academy logo are registered trademarks of the American Academy of Ophthalmology.